

# Far-red fluorescent tags for protein imaging in living tissues

Dmitry SHCHERBO\*, Christopher S. MURPHY†, Galina V. ERMAKOVA\*, Elena A. SOLOVIEVA\*, Tatiana V. CHEPURNYKH‡, Aleksandr S. SHCHEGLOV\*, Vladislav V. VERKHUSHA§, Vladimir Z. PLETNEV\*, Kristin L. HAZELWOOD†, Patrick M. ROCHE†, Sergey LUKYANOV\*<sup>1</sup>, Andrey G. ZARAIKY\*, Michael W. DAVIDSON† and Dmitriy M. CHUDAKOV\*<sup>1,2</sup>

\*Shemiakin-Ovchinnikov Institute of Bioorganic Chemistry, Miklukho-Maklaya 16/10, 117997 Moscow, Russia, †National High Magnetic Field Laboratory and Department of Biological Science, Florida State University, 1800 E. Paul Dirac Dr., Tallahassee, FL 32310, U.S.A., ‡Evrogen JSC, Miklukho-Maklaya 16/10, Moscow 117997, Russia, and §Department of Anatomy and Structural Biology, Albert Einstein College of Medicine, Bronx, NY 10461, U.S.A.

A vast colour palette of monomeric fluorescent proteins has been developed to investigate protein localization, motility and interactions. However, low brightness has remained a problem in far-red variants, which hampers multicolour labelling and whole-body imaging techniques. In the present paper, we report mKate2, a monomeric far-red fluorescent protein that is almost 3-fold brighter than the previously reported mKate and is 10-fold brighter than mPlum. The high-brightness, far-red emission spectrum, excellent pH resistance and photostability, coupled with low toxicity demonstrated in transgenic *Xenopus laevis* embryos, make mKate2 a superior fluorescent tag for imaging in living

tissues. We also report tdKatushka2, a tandem far-red tag that performs well in fusions, provides 4-fold brighter near-IR fluorescence compared with mRaspberry or mCherry, and is 20-fold brighter than mPlum. Together, monomeric mKate2 and pseudo-monomeric tdKatushka2 represent the next generation of extra-bright far-red fluorescent probes offering novel possibilities for fluorescent imaging of proteins in living cells and animals.

**Key words:** far-red fluorescence, fluorescent protein, fusion protein, photostability, whole-body imaging.

## INTRODUCTION

Among the numerous homologues of avGFP (*Aequorea victoria* green fluorescent protein), naturally occurring orange and red fluorescent proteins isolated from other species are either dimeric or tetrameric. For this reason, they are unsuitable in applications as fusion tags for studying localization, intermolecular binding and motility dynamics of proteins of interest [1–3]. To overcome this limitation, monomeric orange, red and far-red fluorescent proteins have been developed in recent years, including the entire palette of ‘mFruits’ [4–6], mKO [7], TagRFP [8] and mKate [9]. However, brightness remains a problem in the far-red monomeric fluorescent proteins. The brightest variant yet reported, mKate, is still dramatically dimmer than any of the wild-type red fluorescent proteins [9]. At the same time, bright far-red monomers are required both for the multicolour labelling in cells and for the whole-body imaging of fusion proteins in living transgenic organisms. In fact, bright far-red fluorescence is a vital characteristic, since the near-IR optical window is exceptionally advantageous for light penetration into living tissues [9]. The present study focused on developing the photophysical characteristics of mKate and has resulted in a new derivative exhibiting dramatic improvements.

## EXPERIMENTAL

Methods concerning cloning and gene construction and characterization of fluorescent proteins *in vitro* can be found in the Supplementary Online Data at <http://www.BiochemJ.org/bj/418/bj4180567add.htm>.

## Microscopy and live-cell imaging

All filters for fluorescence screening and imaging were purchased from Chroma Technology, Omega Filters and Semrock. HeLa epithelial (CCL-2, A.T.C.C., Manassas, VA, U.S.A.) and grey fox lung fibroblast (CCL-168, A.T.C.C.) cells were grown in a 50:50 mixture of DMEM (Dulbecco’s modified Eagle’s medium) and Ham’s F12 with 12.5 % Cosmic calf serum (Hyclone) and transfected with Effectene (Qiagen). Imaging was performed in Delta-T culture chambers (Bioptechs) under a humidified atmosphere of 5 % CO<sub>2</sub> in air. Imaging in widefield mode was performed with a Nikon TE-2000 inverted microscope equipped with Omega QuantaMax™ filters and a Photometrics Cascade II camera or an Olympus IX71 inverted microscope equipped with Semrock BrightLine™ filters and a Hamamatsu ImagEM™ camera. Laser-scanning confocal microscopy was conducted using a Nikon C1Si and an Olympus FV1000, both equipped with argon-ion (457 and 488 nm) and helium–neon or diode (543 and 561 nm respectively) lasers and proprietary filter sets. Spinning disk confocal microscopy was performed on an Olympus DSU-IX81 equipped with a Lumen 200 illuminator (Prior Scientific), a Hamamatsu 9100-13 EMCCD camera, Semrock filters and ten-position filter wheels driven by a Lambda 10-3 controller (Sutter). In some cases, cell cultures expressing fluorescent protein fusions were fixed after imaging in 2 % (w/v) paraformaldehyde (Electron Microscopy Sciences) and washed several times in PBS containing 0.05 M glycine before mounting with a poly(vinyl alcohol)-based medium. Morphological features in all fusion constructs were confirmed by imaging fixed cell preparations on coverslips using a Nikon 80i upright microscope and Semrock

Abbreviations used: avGFP, *Aequorea victoria* green fluorescent protein; CMV, cytomegalovirus; EGFP, enhanced green fluorescent protein; NA, numerical aperture.

<sup>1</sup> Sergey Lukyanov and Dmitriy Chudakov have interest in Evrogen JSC. mKate2, Katushka2 and tdKatushka2 are the property of Evrogen JSC.

<sup>2</sup> To whom correspondence should be addressed (email ChudakovDM@mail.ru).

Texas Red filter set (4040B) coupled to a Hamamatsu Orca ER or a Photometrics CoolSNAP™ HQ² camera.

### Photobleaching assays

Widefield and laser-scanning confocal microscopy photobleaching experiments were conducted with N-terminal fusions of the appropriate fluorescent protein to human histone H2B (six-residue linker) to confine fluorescence to the nucleus in order to closely approximate the dimensions of aqueous droplets of purified fluorescent proteins used in widefield measurements by the Roger Y. Tsien laboratory [1]. HeLa-S3 cells (A.T.C.C.) of average nucleus diameter 17  $\mu\text{m}$  were transfected with the H2B construct using Effectene and maintained in a 5%  $\text{CO}_2$  atmosphere in Bioprotech Delta-T imaging chambers for at least 36 h before imaging. The chambers were transferred to a Bioprotech stage adapter, imaged at low magnification to ensure cell viability, and then photobleached using a 40 $\times$  oil-immersion objective for confocal [Olympus UPlan Apo, NA (numerical aperture) = 1.00] or a 40 $\times$  dry objective (Nikon Plan Fluorite, NA = 0.85) for widefield studies. A metal halide illumination source (EXFO) was used without neutral-density filters for widefield photobleaching assays. Output power at the objective focal plane was measured with a Newport 1918-C meter. Laser lines (543 nm, helium–neon and 488 nm, argon-ion) were adjusted to an output power of 100  $\mu\text{W}$ , measured with a FieldMaxII-TO (Coherent) power meter equipped with a high-sensitivity silicon/germanium optical sensor (Coherent OP-2Vis). The instrument (Olympus FV1000) was set to a zoom of 4 $\times$ , a region of interest of 341.2  $\mu\text{m}^2$  (108 pixels  $\times$  108 pixels), a photomultiplier voltage of 650 V and an offset of 9% with a scan time of 0.181 s/frame. Nuclei having approximately the same dimensions and intensity under the fixed instrument settings were chosen for photobleaching assays. Fluorescence using the 543 nm laser was recorded with a 570 nm dichromatic mirror and 656 nm longpass barrier filter, whereas emission using the 488 nm laser was reflected directly by a mirror through a 510 nm longpass barrier filter. The photobleaching half-times were calculated as described in [6] as the time required to reduce the scan-averaged emission rate to 50% from an initial emission rate of 1000 photons/s per fluorescent protein chromophore.

### Transgenic *Xenopus laevis* embryos

To generate a construct expressing mKate2 under the control of *Xanfl* promoter, the 2200 bp fragment of this promoter sufficient for specifically targeting the expression to the anterior neural plate [10] was subcloned into *VspI* (blunted)/*AgeI* sites of mKate2-N vector, instead of the multiple cloning site region and CMV (cytomegalovirus) promoter. For transgenic experiments, vectors were linearized by *SfiI* and purified using Qiagen columns. Transgenic embryos were generated by the REMI (restriction enzyme-mediated integration) technique exactly as described previously [10] and were analysed using a Leica MZFLIII fluorescent stereomicroscope with the rhodamine filter set: excitation filter 546/10 nm; suppression filter 565 nm. To visualize a fluorescent signal on tissue sections, transgenic embryos were fixed overnight in 4% (w/v) formaldehyde in 0.1% PBS, embedded in 3% agarose in 0.1% PBS and sectioned using a HB 650 Vibratome (Microm) in 50  $\mu\text{m}$  sections. The animal work was performed in Shemyakin and Ovchinnikov Institute of Bioorganic Chemistry, Russian Academy of Sciences, in accordance with the regulations of the Department of Health and Human Services, National Institutes of Health.

## RESULTS AND DISCUSSION

Based on analysis of the mKate crystal structure solved at several pH values (V. Z. Pletnev, unpublished work), we have concluded that the chromophore is capable of forming either a *cis* or *trans* conformation. The *cis* conformation is brightly fluorescent and this increases proportionally with pH, whereas the *trans* conformation is much dimmer and decreases with increasing pH. These data correlate very well with the kindling-like [11] photoactivation behaviour of mKate [9]. Indeed, at physiological pH, mKate exhibits a 10–20% increase in fluorescent brightness in response to irradiation by green light (see [9] and below), which corresponds well to a light-induced chromophore *trans*–*cis* isomerization. Whereas Ser<sup>148</sup> (numbering in accordance with avGFP alignment, Figure 1) stabilizes the fluorescent *cis* conformation of mKate (similar to most red fluorescent proteins, including DsRed [12,13], and kindling fluorescent protein [11,14]), the *trans* conformation of the chromophore is stabilized by Ser<sup>165</sup> (similar to eqFP611 [15] and chromoproteins [16]). We therefore assumed that mutagenesis at position 165 could disrupt the hydrogen-bonding system that stabilizes the chromophore in *trans* conformation, and thus produce a new variant with higher brightness and lower pH-dependence.

Accordingly, we performed saturation mutagenesis (using all 20 amino acids) of Ser<sup>165</sup> and screened for brighter variants of mKate. The brightest and fastest maturing variant at 37°C contained the S165A substitution. This result generally confirmed our assumption: in contrast with the polar hydrophilic residue serine, the non-polar hydrophobic alanine residue is unable to take part in the hydrogen-bonding system that stabilizes the low-fluorescent *trans* conformation of the chromophore. The S165A mutation should therefore increase the portion of the brightly fluorescent *cis* chromophore in the physiological and acidic pH range.

We optimized further the new variant using two rounds of random mutagenesis (see the Supplementary Online Data at <http://www.BiochemJ.org/bj/418/bj4180567add.htm>) and obtained the beneficial mutation V48A, which essentially increases the protein maturation rate, assessed by the relative brightness of the freshly grown *Escherichia coli* expressing mKate variants (results not shown). Additionally, we uncovered a second mutation, K238R, which is found in the wild-type eqFP578 sequence [8] and also seems to enhance maturation slightly. Valine was inserted as the second amino acid in order to generate the optimal Kozak sequence for eukaryotic translation [17]. Collectively, these major and minor changes resulted in a much brighter, faster maturing, acid-resistant and more photostable monomeric far-red fluorescent protein that we named mKate2.

The main characteristics of mKate2 are summarized in Table 1. The spectral profiles are almost identical with those of mKate (Figure 2a), with excitation and emission peaking at 588 and 633 nm respectively. Compared with mKate, the fluorescence of mKate2 is far more pH-stable (Figure 2b), thus increasing its brightness at physiological pH and making mKate2 a more reliable reporter for imaging in acidic organelles, such as late and recycling endosomes and lysosomes. mKate2 is characterized by a ~2-fold higher molar absorption coefficient and 1.4-fold higher fluorescence quantum yield compared with mKate, resulting in 2.8-fold brighter fluorescence. The calculated brightness for mKate2 is approx. 74% of EGFP (enhanced green fluorescent protein), which makes the new variant 10-fold brighter than mPlum (Table 1). Within the optical window optimal for light penetration in living tissues, calculated brightness of mKate2 is at least 2-fold higher compared with any monomeric fluorescent protein reported to date (Figure 2c and Table 1).

**Table 1** Key characteristics of tdKatushka2 and mKate2 in comparison with other monomeric far-red fluorescent proteins

Brightness is calculated as a product of molar absorption coefficient and the fluorescence quantum yield, and is presented as a percentage of the brightness of mEGFP. Photostability is given in s as the normalized (from 1000 photons/molecule/s) fluorescent state half-life upon irradiation. Brightness beyond 650 nm (see Figure 2c) is presented as a percentage of the brightness of tdKatushka2.


Fluorescent protein	$\lambda_{ex}$ (nm)	$\lambda_{em}$ (nm)	Quantum yield	Molar absorption coefficient ( $M^{-1} \cdot cm^{-1}$ )	Brightness (% relative to mEGFP)	Brightness beyond 650 nm (% relative to tdKatushka2)	$pK_a$	Photostability, widefield	Photostability, confocal	Reference(s)
mCherry	587	610	0.22	72 000	47	24	< 4.5	135	2770	[1]
mRaspberry	598	625	0.15	86 000	38	26	—	35	1010	[5,9]
mPlum	590	649	0.10	41 000*	12‡	11‡	< 4.5	77‡	770‡	[5,9]
				22 000†	7§	6§		42§	410§	
mKate	588	635	0.28	31 500	26	19	6.2	71	1260	[9]
mKate2	588	633	0.40	62 500	74	52	5.4	118	2690	—
tdKatushka2	588	633	0.37	66 250 × 2	146	100	5.4	144	4090	—

\*Literature data.


†Our data.

‡Calculated using literature data.

§Calculated using our data.

**Figure 1** Alignment of the amino acid sequences for the selected fluorescent proteins

Structurally important regions are highlighted in grey,  $\beta$ -strands are shown by arrows, and  $\alpha$ -helices are shown by ribbons. Buried residues are shaded. Amino acid residues responsible for the far-red fluorescence of the *cis*-chromophore of Katushka, mKate and mKate2 are underlined. Random mutations generated in the course of mKate optimization are shown white on black, targeted mutations are shown bold. Overall alignment numbering follows that of avGFP.



**Figure 2** Spectral characteristics of mKate2 in comparison with selected fluorescent proteins


(a) Normalized mKate2 absorption (solid line), fluorescence excitation (---) and emission (---) spectra. (b) pH stability of mKate2 (solid line) and mKate (broken line) fluorescence. (c) Emission spectra of mCherry, far-red monomeric fluorescent proteins and tdKatushka2 relative to their calculated brightness (Table 1). Scaling was applied to the area of the peak. The favourable 'optical window' is shaded grey. (d) Normalized photobleaching curves for mCherry, far-red monomeric fluorescent proteins and tdKatushka2 using laser-scanning confocal microscopy. (e) Normalized photobleaching curves, widefield fluorescence microscopy under metal halide illumination.

Importantly, molar absorption coefficients and fluorescence quantum yields for mKate, mKate2 and tdKatushka2 (see below) were measured in direct comparison with mCherry at the physiological pH of 7.5, to allow relevant comparison of theoretical brightness in cytosol of eukaryotic cells. Previously, we reported higher molar absorption coefficient and quantum yield for mKate, measured at pH of 8.2 [9]. However, since mKate is rather pH-sensitive, the pH change from 8.2 to 7.5 leads to pronounced decrease of the protein brightness. In the present paper, we publish corrected data for mKate at pH 7.5.

Our experiments indicate that mKate2 is characterized by fast chromophore maturation at 37 °C. Standard maturation assay (see the Supplementary Online Data at <http://www.BiochemJ.org/bj418/bj4180567add.htm> and [8]) could barely capture the very end of the maturation curve, thus maturation half-time for mKate2 should be <20 min (compared with 40 min for mCherry, 75 min for mKate and 100 min for TagRFP [8,9]). Maturation of the mKate2 red chromophore is complete, as is evident by a single narrow peak in the absorbance spectrum, which is largely identical with the fluorescence excitation spectrum (Figure 2a). We verified

that mKate2 remains monomeric by gel filtration (see the Supplementary Online Data at <http://www.BiochemJ.org/bj418/bj4180567add.htm>).

A high level of photostability is one of the primary characteristics of fluorescent proteins that is necessary for long-term time-lapse imaging. To determine the photostability of mKate2 in comparison with several well-established fluorescent proteins, we performed photobleaching experiments in living HeLa cells using procedures described previously [6]. Photobleaching assays were conducted with selected fluorescent proteins fused to human histone H2B (Figure 3r) using widefield fluorescence microscopy under metal halide illumination with commercial filter sets and laser-scanning confocal microscopy using the 488 and 543 nm spectral lines from an argon-ion and helium-neon laser respectively (see the Experimental section). The raw photobleaching data were normalized to 1000 photons/molecule/s as described previously [1,6] and are listed in Table 1. Remarkably, mKate2 is more photostable under both widefield and confocal illumination than other monomeric far-red proteins, including mKate, mRaspberry and mPlum (Figures 2d and 2e



**Figure 3** Fluorescence imaging of mKate2 fusion vectors


(a)–(g) C-terminal mKate2 fusion constructs: (a) mKate2- $\beta$ -actin-7 (human); (b) mKate2-endosomes-14 (human RhoB GTPase and c-Myc epitope tag); (c) mKate2-lamin B1-10 (human); (d) mKate2-clathrin light chain-15 (human); (e) mKate2-peroxisomes-2 (peroxisomal targeting signal 1; PTS1); (f) mKate2-membrane-5 (20-amino-acid farnesylation signal from c-Ha-Ras); (g) mKate2- $\alpha$ -tubulin-6 (human); (h) mKate2-VASP-5 (mouse vasodilator-stimulated phosphoprotein); (i) mKate2-annexin (A4)-12 (human). (j)–(t) N-terminal fusion constructs: (j) mKate2-paxillin-22 (chicken); (k) mKate2- $\alpha$ -actinin-19 (human non-muscle); (l) mKate2-vimentin-7 (human); (m) mKate2-Golgi-7 (N-terminal 81 amino acids of human  $\beta$ -1,4-galactosyltransferase); (n) mKate2-EB3-7 (human microtubule-associated protein; RP/EB family); (o) mKate2-keratin-17 (human cyokeratin 18); (p) mKate2-zyxin-7 (human); (q) mKate2-Cx43-7 (rat  $\alpha$ -1 connexin 43); (r) mKate2-H2B-6 (human; prophase); (s) mKate2-mitochondria-7 (human cytochrome c oxidase subunit VIII); (t) mKate2-lysosomes-20 (human lysosomal membrane glycoprotein 1). The cell line used for expressing mKate2 fusion vectors was grey fox lung fibroblast cells (FoLu) in panels (a), (g) and (n), and human cervical adenocarcinoma cells (HeLa; ATCC, CCL2) in all other panels. A different strain of HeLa (ATCC, S3) was used for mitosis studies in (r). Scale bars, 10  $\mu$ m. For each fusion protein, the linker amino acid length is indicated after the name of the targeted organelle or fusion protein.

and Table 1). In contrast with mKate, mKate2 demonstrates almost no kindling (photoactivation) phenomena, consistent with the structural considerations discussed above. We presume that, at physiological pH, the chromophore of mKate2 exists mostly as the *cis* isomer, with limited potential for additional light-induced *trans*–*cis* isomerization of the chromophore to a bright *cis* conformation.

To characterize mKate2 in living cells, we performed transient transfections of HeLa and Phoenix Eco cell lines with a vector driving mKate2 expression without any fusion or localization signals. The fluorescent signal was evenly distributed within the cytosol and nucleus with no visible aggregates or non-specific

localization observed within 4 days of transfection in both cell lines. These data indicate that mKate2 should be non-toxic when expressed in stable cell lines, as well as the previously reported mKate (cell lines available from Marinpharm GmbH).

To verify the utility of mKate2 as a fusion tag, we constructed 20 vectors and expressed them in epithelial and fibroblast cell lines to examine performance. mKate2 performed well in most C- and N-terminal fusions (Figure 3), similar to the results observed with the parent proteins, mKate and TagRFP, and demonstrated excellent brightness and photostability that allowed the capture of a long time series (see Supplementary Videos S1–S7 at <http://www.BiochemJ.org/bj/418/bj4180567add.htm>). A fusion



**Figure 4** Imaging of mKate2 in *X. laevis* embryos

(A) Expression of mKate2 under the *Xanfl* promoter in the transgenic embryos at stage 28 is specifically localized in the forehead region, including eyes, the forebrain and nasal placodes. The embryo is shown from the right-hand side, with the dorsal side at the top and left. (B) Control images, non-transgenic embryo at the same stage. (C) Upper image: expression of mKate2–zyxin under CMV promoter in the transgenic embryo of stage 35 (2 days). Despite quite intensive and ubiquitous expression of mKate2–zyxin, the embryo appears normal and healthy. Lower image: control embryo of the same stage. The embryos are shown from the right-hand side, with the dorsal side at the top.

of mKate2 to annexin (A4) translocated as expected from the cytoplasm to the nuclear and plasma membranes upon induction with ionomycin (Figure 3i and Supplementary Video S2). mKate2 also performed well in fusions targeting the Golgi complex (Figure 3m), clathrin vesicles (Figure 3d), peroxisomes (Figure 3e), lysosomes (Figure 3t), mitochondria (Figure 3s), endosomes (Figure 3b) and gap junctions (Figure 3q), as well as assorted nuclear, focal adhesion and cytoskeletal targets (Figures 3c, 3g, 3h, 3k, 3l, 3n, 3o and 3p).

To verify the low toxicity of mKate2 in transgenic animals, we expressed mKate2 under the control of *Xanfl* promoter in *X. laevis* embryos. We have used the 2200 bp fragment of this promoter, which is sufficient for specific targeting of the expression to the anterior neural plate [10]. As expected, we observed bright red fluorescence in the forehead region, including eyes, the forebrain and nasal placodes (Figures 4A and 4B).


To compare brightness and maturation rates of mKate and mKate2 *in vivo*, *X. laevis* embryos were microinjected with mKate-N and mKate2-N vectors at the stage of two blastomeres. Living embryos were photographed from the animal pole side at the middle gastrula stages (10.5 h after fertilization). As expected, embryos microinjected with mKate2-N demonstrated superior brightness (see Supplementary Figure S1 at <http://www.BiochemJ.org/bj/418/bj4180567add.htm>).

In addition, to test in an embryonic model the performance of mKate2 in a targeting protein fusion, we generated transgenic *X.*

*laevis* embryos bearing a CMV–mKate2–zyxin fusion construct. Despite quite extensive and ubiquitous expression of mKate2–zyxin under the control of the CMV promoter, these embryos appear normal and healthy (Figure 4C and Supplementary Figures S2 and S3 at <http://www.BiochemJ.org/bj/418/bj4180567add.htm>). Importantly, the tadpoles bearing mKate2 and mKate2–zyxin transgenic constructs have demonstrated roughly comparable viability with tadpoles expressing EGFP transgenes (on average 39 % of transgenes selected at stage 46 reached stage 60), which was observed in our previous experiments ([10,18] and A. G. Zaraksky, unpublished work). This indicates that mKate2, as well as EGFP, exerts a low toxic effect on living cells in transgenic organisms.

The dimeric far-red fluorescent protein Katushka is a close relative of mKate and mKate2 [9]. It was therefore intriguing to verify the influence of the key S165A substitution on the spectral properties of Katushka. As could be expected, site-directed mutagenesis S165A resulted in a notable (approx. 20 %) increase of fluorescence brightness. Similarly to mKate2, Katushka2 lost the kindling property and mostly preserved the spectral characteristics of the parental Katushka. To generate a Katushka2 variant suitable for fusions, we employed the popular ‘tandem’ approach [4,19] to create a pseudo-monomeric probe. In spite of the larger size, tandem versions of dimeric fluorescent proteins behave similarly to the true monomers and usually perform well in fusions with many proteins of interest. Indeed, the resulting





**Figure 5** Fluorescence imaging of tdKatushka2 fusion vectors

(a)–(g) C-terminal tdKatushka2 fusion constructs: (a) tdKatushka2–clathrin light chain-15 (human); (b) tdKatushka2– $\beta$ -actin-7 (human); (c) tdKatushka2–VASP-5 (mouse vasodilator-stimulated phosphoprotein); (d) tdKatushka2–lamin B1-10 (human); (e) tdKatushka2–membrane-5 (20-amino acid farnesylation signal from c-Ha-Ras); (f) tdKatushka2–fibrillarin-7 (human); (g) tdKatushka2–peroxisomes-2 (peroxisomal targeting signal 1; PTS1). (h)–(p) N-terminal tdKatushka2 fusion constructs: (h) tdKatushka2–EB3-7 (human microtubule-associated protein; RP/EB family); (i) tdKatushka2–Golgi-7 (N-terminal 81 amino acids of human  $\beta$ -1,4-galactosyltransferase); (j) tdKatushka2–zyxin-7 (human); (k) tdKatushka2–Cx26-7 (rat  $\beta$ 2 connexin 26); (l) tdKatushka2–paxillin-22 (chicken); (m) tdKatushka2–lysosomes-20 (human lysosomal membrane glycoprotein 1); (n) tdKatushka2–VE-cadherin (human vascular epithelial cadherin); (o) tdKatushka2–VSVG-12 (vesicular stomatitis virus glycoprotein; membrane); (p) tdKatushka2–CENP-B22 (human CENP-B DNA-binding domain). The cell line used for expressing tdKatushka2 fusion vectors was grey fox lung fibroblast cells (FoLu) in panel (h) and human cervical adenocarcinoma cells (HeLa; ATCC, CCL2) in all other panels. Scale bars, 10  $\mu$ m. For each fusion protein, the linker amino acid length is indicated after the name of the targeted organelle or fusion protein.

tdKatushka2 performed well in a number of fusions, including those targeting clathrin vesicles, the nuclear lamina, nucleoli, the Golgi, peroxisomes, centrosomes, lysosomes, plasma membrane, focal adhesions and even several fusion proteins that themselves form biopolymers, including actin and gap junctions (Figure 5). Further characterization *in vitro* and in living cells showed that extra-bright (1.46-fold brighter than EGFP per molecule) far-red tdKatushka2 demonstrates fast maturation, high pH-stability and high photostability (Table 1).

Advantageous characteristics of mKate2 and tdKatushka2 make them probably the best fluorescent tags yet reported in the far-red range. Bright monomeric fluorescent proteins now span almost the entire visible range, which greatly expands the possibilities of using these probes as monitors of gene expression, multicolour labelling, dynamic interactions, and FRET (fluorescence resonance energy transfer)/FLIM (fluorescence lifetime imaging microscopy) techniques. Importantly, the emission spectra of mKate2 and tdKatushka2 extend into the near-IR optical window that is optimal for light penetration into living tissues [9]. This unique combination of high brightness and far-red emission definitely makes mKate2 and tdKatushka2 the fluorescent proteins

of choice for visualizing fusion tags of interest in whole-body imaging of live animals.

#### AUTHOR CONTRIBUTIONS

D.S., T.V.C. and A.S.S. generated and analysed mKate variants. K.L.H., P.M.R. and C.S.M. generated fusion constructs and performed microscopy and photobleaching experiments. G.V.E., E.A.S. and A.G.Z. performed experiments with *X. laevis* embryos. V.Z.P. analysed the crystal structure of mKate. V.V.V., S.L., M.W.D. and D.M.C. designed and planned the project and wrote the manuscript.

#### FUNDING

This work was supported by the Howard Hughes Medical Institute [grant numbers 55005618 and 55005630], Molecular and Cell Biology Program, Russian Academy of Sciences, RFBR 08-04-01702-a, European Union Framework Programme 6 Integrated Project [grant number LSHG-CT-2003-503259], 'State Support of the Leading Scientific Schools' programme [grant number NS-2395.2008.4], National Institutes of Health [grant numbers GM070358 and GM073913], Russian Federal Agency for Science and Innovations [grant number 02.513.12.3013]. D.M.C. is supported by the Grant of the President of Russian Federation [grant number MK-6119.2008.4] and the Russian Science Support Foundation.

## REFERENCES

- Shaner, N. C., Steinbach, P. A. and Tsien, R. Y. (2005) A guide to choosing fluorescent proteins. *Nat. Methods* **2**, 905–909.
- Chudakov, D. M., Lukyanov, S. and Lukyanov, K. A. (2005) Fluorescent proteins as a toolkit for *in vivo* imaging. *Trends Biotechnol.* **23**, 605–613.
- Shaner, N. C., Patterson, G. H. and Davidson, M. W. (2007) Advances in fluorescent protein technology. *J. Cell Sci.* **120**, 4247–4260.
- Shaner, N. C., Campbell, R. E., Steinbach, P. A., Giepmans, B. N., Palmer, A. E. and Tsien, R. Y. (2004) Improved monomeric red, orange and yellow fluorescent proteins derived from *Discosoma* sp. red fluorescent protein. *Nat. Biotechnol.* **22**, 1567–1572.
- Wang, L., Jackson, W. C., Steinbach, P. A. and Tsien, R. Y. (2004) Evolution of new nonantibody proteins via iterative somatic hypermutation. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 16745–16749.
- Shaner, N. C., Lin, M. Z., McKeown, M. R., Steinbach, P. A., Hazelwood, K. L., Davidson, M. W. and Tsien, R. Y. (2008) Improving the photostability of bright monomeric orange and red fluorescent proteins. *Nat. Methods* **5**, 545–551.
- Karasawa, S., Araki, T., Nagai, T., Mizuno, H. and Miyawaki, A. (2004) Cyan-emitting and orange-emitting fluorescent proteins as a donor/acceptor pair for fluorescence resonance energy transfer. *Biochem. J.* **381**, 307–312.
- Merzlyak, E. M., Goedhart, J., Shcherbo, D., Bulina, M. E., Shcheglov, A. S., Fradkov, A. F., Gaintzeva, A., Lukyanov, K. A., Lukyanov, S., Gadella, T. W. and Chudakov, D. M. (2007) Bright monomeric red fluorescent protein with an extended fluorescence lifetime. *Nat. Methods* **4**, 555–557.
- Shcherbo, D., Merzlyak, E. M., Chepurnykh, T. V., Fradkov, A. F., Ermakova, G. V., Solovieva, E. A., Lukyanov, K. A., Bogdanova, E. A., Zaraisky, A. G., Lukyanov, S. and Chudakov, D. M. (2007) Bright far-red fluorescent protein for whole-body imaging. *Nat. Methods* **4**, 741–746.
- Martynova, N., Eroshkin, F., Ermakova, G., Bayramov, A., Gray, J., Grainger, R. and Zaraisky, A. (2004) Patterning the forebrain: FoxA4/Pintallavis and Xvent2 determine the posterior limit of Xanf1 expression in the neural plate. *Development* **131**, 2329–2338.
- Chudakov, D. M., Feofanov, A. V., Mudrik, N. N., Lukyanov, S. and Lukyanov, K. A. (2003) Chromophore environment provides clue to “kindling fluorescent protein” riddle. *J. Biol. Chem.* **278**, 7215–7219.
- Wall, M. A., Socolich, M. and Ranganathan, R. (2000) The structural basis for red fluorescence in the tetrameric GFP homolog DsRed. *Nat. Struct. Biol.* **7**, 1133–1138.
- Yarbrough, D., Wachter, R. M., Kallio, K., Matz, M. V. and Remington, S. J. (2001) Refined crystal structure of DsRed, a red fluorescent protein from coral, at 2.0-Å resolution. *Proc. Natl. Acad. Sci. U.S.A.* **98**, 462–467.
- Quillin, M. L., Anstrom, D. M., Shu, X., O’Leary, S., Kallio, K., Chudakov, D. M. and Remington, S. J. (2005) Kindling fluorescent protein from *Anemonia sulcata*: dark-state structure at 1.38 Å resolution. *Biochemistry* **44**, 5774–5787.
- Petersen, J., Wilmann, P. G., Beddoe, T., Oakley, A. J., Devenish, R. J., Prescott, M. and Rossjohn, J. (2003) The 2.0-Å crystal structure of eqFP611, a far red fluorescent protein from the sea anemone *Entacmaea quadricolor*. *J. Biol. Chem.* **278**, 44626–44631.
- Beddoe, T., Ling, M., Dove, S., Hoegh-Guldberg, O., Devenish, R. J., Prescott, M. and Rossjohn, J. (2003) The production, purification and crystallization of a pocilloporin pigment from a reef-forming coral. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **59**, 597–599.
- Kozak, M. (1984) Point mutations close to the AUG initiator codon affect the efficiency of translation of rat preproinsulin *in vivo*. *Nature* **308**, 241–246.
- Ermakova, G. V., Solovieva, E. A., Martynova, N. Y. and Zaraisky, A. G. (2007) The homeodomain factor Xanf represses expression of genes in the presumptive rostral forebrain that specify more caudal brain regions. *Dev. Biol.* **307**, 483–497.
- Fradkov, A. F., Verkhusha, V. V., Staroverov, D. B., Bulina, M. E., Yanushevich, Y. G., Martynov, V. I., Lukyanov, S. and Lukyanov, K. A. (2002) Far-red fluorescent tag for protein labelling. *Biochem. J.* **368**, 17–21.

Received 26 September 2008/1 December 2008; accepted 4 December 2008

Published as BJ Immediate Publication 15 January 2009, doi:10.1042/BJ20081949




## SUPPLEMENTARY ONLINE DATA

# Far-red fluorescent tags for protein imaging in living tissues

Dmitry SHCHERBO\*, Christopher S. MURPHY†, Galina V. ERMAKOVA\*, Elena A. SOLOVIEVA\*, Tatiana V. CHEPURNYKH‡, Aleksandr S. SHCHEGLOV\*, Vladislav V. VERKHUSHA§, Vladimir Z. PLETNEV\*, Kristin L. HAZELWOOD†, Patrick M. ROCHE†, Sergey LUKYANOV\*<sup>1</sup>, Andrey G. ZARAIKY\*, Michael W. DAVIDSON† and Dmitriy M. CHUDAKOV\*<sup>1,2</sup>

\*Shemiakin-Ovchinnikov Institute of Bioorganic Chemistry, Miklukho-Maklaya 16/10, 117997 Moscow, Russia, †National High Magnetic Field Laboratory and Department of Biological Science, Florida State University, 1800 E. Paul Dirac Dr., Tallahassee, FL 32310, U.S.A., ‡Evrogen JSC, Miklukho-Maklaya 16/10, Moscow 117997, Russia, and §Department of Anatomy and Structural Biology, Albert Einstein College of Medicine, Bronx, NY 10461, U.S.A.



**Figure S1 Comparison of mKate and mKate2 in developing *X. laevis* embryos**

At the stage of two blastomeres, embryos were microinjected with mKate-N1 and mKate2-N vectors. Living embryos were photographed from the animal pole side at the middle gastrula stages (10.5 h after fertilization).

## EXPERIMENTAL

### Cloning and gene construction

Synthetic DNA oligonucleotides for cloning and mutagenesis were purchased from Integrated DNA Technologies or Evrogen. PCR products and products of restriction digests were purified by gel electrophoresis and extraction using the QIAquick™ gel extraction kit (Qiagen). Plasmid DNA was purified from overnight cultures by using the QIAprep Spin Miniprep kit (Qiagen). Restriction endonucleases were purchased from Invitrogen or New England Biolabs. Sequencing was used to confirm the complete cDNA sequences for all fluorescent protein variants and fusion constructs (Florida State University Bioanalytical and Molecular Cloning DNA Sequencing Laboratory or Evrogen). Site-directed mutagenesis was performed by overlap-extension PCR [1] with primers containing the appropriate target substitutions. The Diversity PCR Random Mutagenesis kit (Clontech) was used for random mutagenesis in conditions optimal for seven mutations per 1000 bp. For bacterial expression, a PCR-amplified BamHI/HindIII fragment encoding a fluorescent protein variant was cloned into the pQE30 vector (Qiagen).


For expression in eukaryotic cells, a PCR-amplified AgeI/NotI fragment encoding a fluorescent protein variant was swapped for TurboGFP within the pTurboGFP-N or pTurboGFP-C vectors (Evrogen). To generate fusion vectors, the appropriate cloning vector and an EGFP fusion vector were digested, either sequentially or doubly, with the appropriate enzymes and ligated together after gel purification. Thus, to prepare mKate2 N-terminal fusions, the following digests were performed: human non-muscle  $\alpha$ -actinin, EcoRI and NotI (vector source, Tom Keller, Florida State University); human cytochrome *c* oxidase subunit VIII, BamHI and NotI (mitochondria, Clontech); human zyxin, BamHI and NotI (Clare Waterman-Storer, National Institutes of Health, Bethesda, MD, U.S.A.); rat  $\alpha$ -1 connexin-43, EcoRI and BamHI (Matthias Falk, Lehigh University, Bethlehem, PA, U.S.A.); human H2B, BamHI and NotI (George Patterson, National Institutes of Health); N-terminal 81 amino acids of human  $\beta$ -1,4-galactosyltransferase, BamHI and NotI (Golgi, Clontech); human microtubule-associated protein EB3, BamHI and NotI (Lynne Cassimeris, Lehigh University); human vimentin, BamHI and NotI (Robert Goldman, Northwestern University Medical School, Chicago, IL, U.S.A.); human keratin 18, EcoRI and NotI (Open Biosystems); chicken paxillin, EcoRI and NotI (Alan Horwitz, University of Virginia, Charlottesville, VA, U.S.A.); and human lysosomal membrane glycoprotein 1, BamHI and NotI (LAMP1; George Patterson). To prepare mKate2 C-terminal fusions, the following digests were performed: human  $\beta$ -actin, NheI and BglII (Clontech); human  $\alpha$ -tubulin, NheI and BglII (Clontech); human light chain clathrin, NheI and BglII (George Patterson); human lamin B1, NheI and BglII (George Patterson); PTS1 (peroxisomes), AgeI and BspEI (Clontech); human annexin (A4), AgeI and BspEI (Alen Piljic, EMBL, Heidelberg, Germany); human RhoB GTPase with an N-terminal c-Myc epitope tag (endosomes), AgeI and BspEI (Clontech); and the 20-amino-acid farnesylation signal from c-Ha-Ras, AgeI and BspEI (membrane, Clontech); mouse VASP, NheI and BglII (Robert Fischer, National Institutes of Health). DNA for mammalian transfection was prepared using either the Plasmid Midi or Maxi kit (Qiagen).

### Characterization of fluorescent proteins *in vitro*

Proteins fused to the N-terminal polyhistidine tag were expressed in *E. coli* XL1 Blue strain (Invitrogen). The bacterial cultures were centrifuged at 4000 *g* for 20 min, and the cell pellets were resuspended in 20 mM Tris/HCl (pH 7.5) and 100 mM NaCl and lysed by sonication. The recombinant proteins were purified using TALON™ metal-affinity resin (Clontech) followed by a desalting step over gel-filtration columns (Bio-Rad Laboratory).


<sup>1</sup> Sergey Lukyanov and Dmitriy Chudakov have interest in Evrogen JSC. mKate2, Katushka2 and tdKatushka2 are the property of Evrogen JSC.

<sup>2</sup> To whom correspondence should be addressed (email ChudakovDM@mail.ru).



**Figure S2** Expression of mKate2-zyxin in *X. laevis* embryos

Expression of mKate2-zyxin under CMV promoter control in the transgenic *X. laevis* embryo of stage 47–48 (7 days). Despite quite intensive and ubiquitous expression of mKate2-zyxin, the embryos look normal and healthy. The embryo is shown from the left-hand side, with the dorsal side at the top.



**Figure S3** Expression of mKate2-zyxin in epidermal cells of *X. laevis* embryo

The epidermal cells of stage 35 transgenic embryo, expressing mKate2-zyxin under the control of the CMV promoter, as it is seen on the transverse section of the embryo. mKate2-zyxin is localized primarily in the submembrane space of the epidermal cells, where it presumably interacts with the membrane-associated cytoskeleton proteins.

Absorption spectra were recorded with a Beckman DU520 UV–visible spectrophotometer. A Varian Cary Eclipse fluorescence spectrophotometer was used for measuring excitation–emission spectra. All measurements were performed in 20 mM Tris/HCl (pH 7.5) and 100 mM NaCl. For molar absorption coefficient determination, we relied on measuring mature chromophore concentration. Proteins were alkali-denatured with an equal volume of 2 M NaOH, and absorbance spectra were measured immediately. It is known that alkali-denatured DsRed-like chromophore converts into the GFP (green fluorescent protein)-like one [2], with an absorption coefficient of  $44\,000\text{ M}^{-1}\cdot\text{cm}^{-1}$  at 452 nm under these conditions. Based on the absorption of the

native and alkali-denatured proteins, molar absorption coefficients for the native states were calculated. For determination of the quantum yield, the fluorescence of the mutant variants was directly compared with that of mCherry (quantum yield 0.22), and corrected to the area of the peak. pH titrations were performed by using a series of buffers in the range 3–10. For each pH value, an aliquot of purified protein was diluted in an equal volume of the corresponding buffer solution, and the fluorescence brightness was measured after 1 h of incubation at room temperature (25 °C). For accuracy, the actual pH was measured in each sample using a microelectrode (Sartorius). Gel filtration was performed on a 1 cm × 90 cm Econo-column (Bio-Rad Laboratories) packed with Sephacryl-S200 (Amersham Biosciences), equilibrated with 25 mM Tris/HCl (pH 7.5) and 150 mM NaCl. To measure the maturation rate, transformed *E. coli* cells (XL1 Blue strain) were grown overnight in LB (Luria–Bertani) supplemented with 200 mg/l ampicillin and 2 % D-glucose. Tubes were filled to the rim and sealed upon 1 h of induction to restrict oxygen availability. The bacterial cultures were centrifuged at 4000 *g* for 15 min, and the cell pellets were resuspended in 20 mM Tris/HCl (pH 7.5) and 100 mM NaCl and lysed by sonication. The recombinant proteins were purified using TALON™ metal-affinity resin (Clontech). Maturation was performed at 37 °C, in 35 mM KCl, 2 mM MgCl<sub>2</sub>, 50 mM Tris/HCl (pH 7.5) and 1 mM dithiothreitol. A Varian Cary Eclipse fluorescence spectrophotometer was used for measuring maturation kinetics.

## REFERENCES

- 1 Ho, S. N., Hunt, H. D., Horton, R. M., Pullen, J. K. and Pease, L. R. (1989) Site-directed mutagenesis by overlap extension using the polymerase chain reaction. *Gene* **77**, 51–59.
- 2 Gross, L. A., Baird, G. S., Hoffman, R. C., Baldridge, K. K. and Tsien, R. Y. (2000) The structure of the chromophore within DsRed, a red fluorescent protein from coral. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 11990–11995.

Received 26 September 2008/1 December 2008; accepted 4 December 2008  
Published as BJ Immediate Publication 15 January 2009, doi:10.1042/BJ20081949