Meier et al. Vol. 127, No. 6, December 1994. Pages 1505-1514.

The underlined and bold portion of the corrected rat NAP57 sequence below represents new carboxy-terminal amino acids due to the insertion of an extra guanoside nucleotide at position 1464 of the NAP57 cDNA.

MADAEAAMTFPKKHKKKKERKPLPEADVAEIQHAEDFLIKPESKAAQLDT	50
SQWPLLLKNFDRLNVRTTHYTPIPCGSNPLKREIGEYVRTGFINLDKPSN	100
PSSHEVVAWIRRILRVEKTGHSGTLDPKVTGCLIVCIERATRLVKSQQSA	150
GKEYVGVVRLHNAIEGTAQLSRALETLTGALFQRPPLIAAVKRQLRVRTI	200
YESRVVEYDPERRLGVFWVSCEAGTYIRTLCVHLGLLLGVGGQMQELRRV	250
RSGVVGERDHMVTMHDVLDAQYLYDHHRDESYLRRVVFPLEKLLTSHKRL	300
VMKDSAVNAICYGAKIMLPGLLRYEDGIEVNQEVVVITTKGEAVCVAIAL	350
MTTAVISTCDHGVVAKIKRVIMERDTYPRKWGLGPKASQKKQLIKQGLLD	400
KHGRPTDGTPASWTRDYVDYSDSSKKATAAEATPGPGVTADAASIVKRKR	450
DSDSDADE <b>ATPTTTPRVKKEKKKKKEKADGGEEAAEDGDGDATRKKKKKK</b>	500
ARAAEELSG	509