

Characterization of Phenotypic Switching in *Cryptococcus neoformans* Biofilms

Luis R. Martinez · David C. Ibom ·
Arturo Casadevall · Bettina C. Fries

Received: 4 March 2008 / Accepted: 13 May 2008 / Published online: 21 June 2008
© Springer Science+Business Media B.V. 2008

Abstract *Cryptococcus neoformans* is an encapsulated yeast-like fungus that is a relatively frequent cause of meningoen- cephalitis in immunocompromised patients and also occasionally causes disease in apparently healthy individuals. This fungus collectively forms biofilms on polystyrene plates and medical devices, whereas individually can undergo phenotypic switching. Both events have profound consequences in the establishment of fungal infection and are associated with persistent infection due to increase resistance to antimicrobial therapy. In this study, we characterized switch phenotypes in *C. neoformans* biofilms. Smooth, mucoid, and wrinkled switch phenotypes of various switching *C. neoformans*

strains were examined for their adhering and biofilm-forming ability on 96-well plates using cell counts and 2,3-bis (2-methoxy-4-nitro-5-sulfophenyl)-5-[(phenylamino) carbonyl]-2H-tetrazolium hydroxide (XTT) reduction assay, respectively. Both assays showed that *C. neoformans* strains with the parent smooth phenotype adhered and formed stronger biofilms than their mucoid and wrinkled counterparts. Furthermore, the phenotypic switching frequencies of the individual colony types grown in biofilms or as planktonic cells were investigated. For the parent smooth variant of most strains, we found enhanced phenotypic switching in cryptococcal biofilms when compared to switching rates of planktonic cells. In contrast, the back-switching rate of mucoid to smooth variant was significantly higher in planktonic cells of seven strains of *C. neoformans* strains. These results suggested that phenotypic switching can occur in cryptococcal biofilms and extend our understanding of the relationship of both phenomena.

L. R. Martinez · B. C. Fries (✉)
Division of Infectious Diseases, Department of Medicine,
Albert Einstein College of Medicine, 1300 Morris Park
Avenue, Bronx, NY 10461, USA
e-mail: fries@aecom.yu.edu

L. R. Martinez · A. Casadevall · B. C. Fries
Department of Microbiology and Immunology, Albert
Einstein College of Medicine, 1300 Morris Park Avenue,
Bronx, NY 10461, USA

D. C. Ibom
Department of Natural Sciences, State University
of New York, Purchase, NY, USA

A. Casadevall
Department of Microbiology, Albert Einstein College of
Medicine, 1300 Morris Park Avenue, Bronx,
NY 10461, USA

Keywords Fungi · Biofilms · Phenotypic switching · Morphology · Planktonic

Introduction

Cryptococcus neoformans is an encapsulated yeast-like fungus that is a relatively frequent cause of meningoen- cephalitis in immunocompromised patients and also

occasionally causes disease in apparently healthy individuals [1]. This fungus has a polysaccharide capsule composed primarily of glucuronoxylomannan (GXM) that is a major contributor to *C. neoformans* virulence, since non-encapsulated strains are not pathogenic [2]. *C. neoformans* enters the host by the respiratory route in the form of dehydrated haploid yeast or basidiospores, which are mostly found in the environment in association with soil contaminated with pigeon excreta [3].

Phenotypic switching occurs in A, B, and D serotypes of *C. neoformans* strains [4] and is emerging as a fundamental mechanism of virulence that may allow persistence of infection by promoting the generation of a new variant that successfully escapes the immune response [5]. *C. neoformans* can undergo phenotypic switching from a smooth parent variant to mucoid, serrated, or wrinkled colony variants with altered virulence [4]. Furthermore, for strain RC-2 switching to a mucoid colony variant during chronic murine infection was documented and associated with poor outcome. In addition, antifungal drug therapy can promote selection of a hypervirulent mucoid variant [6]. This is relevant because frequently treatment failures in immunocompromised hosts are the result of persistence of the initial strain despite adequate antifungal therapy [7].

C. neoformans biofilm formation has profound consequences in the establishment of fungal infection and is associated with persistent infection since biofilms increase resistance to host immune mechanisms [8] and antimicrobial therapy [9]. This fungus can form biofilms on polystyrene plates [10] and medical devices after GXM shedding [11]. A typical cryptococcal biofilm consists of a complex network of yeast cells enmeshed in a substantial amount of polysaccharide matrix [9, 10].

Since phenotypic switching and biofilm formation are two mechanisms associated with persistent *C. neoformans* infection, we hypothesized that biofilm formation may constitute a niche that promotes phenotypic switching and therefore investigated the relationship of these events.

Materials and Methods

Strains

Strain 24067, a serotype D strain (also known as 52D), was originally isolated from a case of human

cryptococcosis and can be obtained from the American Type Culture Collection (Rockville, Md.). Several stable and switching variants of the strain have been identified and are described elsewhere [12]. The variant 24067A arose spontaneously during passage in vitro and was identified as different from its parent strain because it was hypovirulent for mice. Three other smooth parent variants of 24067 F4-SM and WR-SM and RC-2-SM also switch to wrinkled (F4-WR, WR) and mucoid colonies (RC-2 MC, F4-MC). They all arouse spontaneously and reversibly switch. SB4 is a serotype A strain that can switch between SM, WR, and serrated colony types.

Biofilm Formation

C. neoformans cells were collected by centrifugation, washed twice with phosphate-buffered saline (PBS), counted using a hemacytometer, and suspended at 10^7 cells/ml in minimal media (20 mg/ml thiamine, 30 mM glucose, 26 mM glycine, 20 mM $MgSO_4 \cdot 7H_2O$, and 58.8 mM KH_2PO_4). Then, 100 μ l of the suspension was added into individual wells of polystyrene 96-well plates (Fisher, MA) and incubated at 37°C without shaking. Biofilms were formed for 48 h. Following biofilm development, the wells containing *C. neoformans* biofilms were washed three times with PBS to remove non-adhered cryptococcal cells. As described in detail elsewhere fungal cells that remained attached to the plastic surface were considered true biofilms [9].

Measurement of Biofilm Metabolic Activity by XTT-Reduction Assay

A semiquantitative measurement of *C. neoformans* biofilm formation was obtained from the 2,3-bis (2-methoxy-4-nitro-5-sulfophenyl)-5-[(phenylamino) carbonyl]-2H-tetrazolium hydroxide (XTT) reduction assay [13]. For *C. neoformans* strains, 50 μ l of XTT salt solution (1 mg/ml in PBS) and 4 μ l of menadione solution (1 mM in acetone; Sigma) were added to each well. Microtiter plates were incubated at 37°C for 5 h. Fungal mitochondrial dehydrogenases activity reduces XTT tetrazolium salt to XTT formazan resulting in colorimetric change that correlates with cell viability [13]. The colorimetric change was measured using a microtiter reader (Labsystem Multiskan MS, Finland) at 492 nm.

Adhesion Assay

For each *C. neoformans* strain, 10^3 cells were added into individual wells of polystyrene 96-well plates (Fisher, MA) and incubated at 37°C for 2 h. After incubation, wells were washed thrice with PBS to get rid of non-adhered fungal cells. Adhesion was determined by cell counts using a PhotoZoom inverted light microscope (Cambridge Instrument, MA). The number of cryptococcal cells attached to the bottom of each well was averaged per 40-power field. This assay was done in triplicates. Differences in adherence were confirmed by determining CFU of adhered cells after plating on Sabouraud dextrose agar plates.

Comparison of Biofilm and Planktonic Cryptococcal Cell Phenotypic Switching Rate

Fungal Biofilms

C. neoformans biofilms were scraped from the bottom of the wells with a sterile 200 µl micropipette tip to dissociate yeast cells. A volume of 100 µl of suspension containing dissociated cells was aspirated from the wells, transferred to an Eppendorf tube with 900 µl of PBS and vortexed gently for 3 min. Single cell suspension was confirmed by microscopy. To determine switching rate of biofilm-associated cells 100 µl of a 4×10^4 cells/ml suspension were plated on Sabouraud dextrose agar plates and incubated at 30°C for 48 h.

Planktonic Cells

Planktonic cells of individual *C. neoformans* strains and switch variants were grown as above in minimal media for 48 h at 37°C, collected by centrifugation, washed twice with PBS, and counted using a hemacytometer. To determine switching rate of planktonic cells 100 µl of a 4×10^4 cells/ml suspension were plated on Sabouraud dextrose agar plates and incubated at 30°C for 48 h.

Frequencies of switching for the individual cell types were determined by visually scoring colonies with altered morphology. Approximately 40,000 colonies were screened.

Statistical Analysis

All data were subjected to statistical analysis using Origin 7.0 (Origin Lab Corporation, Northampton, MA). *P* values were calculated by *t*-test. *P* values of <0.05 were considered significant.

Results

The Parent Smooth Phenotypes of Switching *C. neoformans* Strains Form Stronger Biofilms than their Mucoïd and Wrinkled Switch Variants

Biofilm formation by the individual switch variants of *C. neoformans* strains RC-2, F4-24067, WRS/SM, SB-4 on polystyrene microtiter plates were compared using the colorimetric XTT-reduction assay. Interestingly, the smooth parent phenotype produced stronger biofilms when compared to the mucoïd or wrinkled switched phenotype strains (Fig. 1a). For instance, RC-2 SM strain formed significantly stronger biofilm than its RC-2 MC counterpart (Fig. 1b).

Enhanced Biofilm Formation of the SM Parents is Associated with Enhanced Adherence of this Phenotype to the Plastic Surface of the Microtiter Plates

Adhesion of cryptococcal cells to the polystyrene surface was examined using cell counts and light microscopy. There was a strong correlation between the adhesion and biofilm formation among all the examined strains. The smooth phenotype of *C. neoformans* strains became firmly attached to the plastic surface of the microtiter plate, whereas the mucoïd and wrinkled switch variant did not attach as well (Fig. 2a). Accordingly adhesion percentages for RC-2 SM (Fig. 2b) and WRS/SM-SM were 80 and 50% higher than RC-2 MC (Fig. 2c) and WRS/SM-MC, respectively.

C. neoformans Biofilms and Planktonic Cells Differed in Phenotypic Switching Frequency

We investigated whether or not biofilms constituted a microenvironment that promotes phenotypic switching to different colony morphology (small, wrinkled, serrated, sectoried, mucoïd). Table 1 summarizes the




Fig. 1 *C. neoformans* phenotype affects biofilm formation. **(a)** Cells from smooth and mucoid phenotypes from four *C. neoformans* isolates were allowed to adhere to polystyrene plates and the amount of biofilm formed after 48 h was determined by using XTT assay. Bars are the averages of three measurements, and error bars denote standard deviations. Asterisks denote *P* value significance calculated by *t*-test.

P values of <0.05 were considered significant. **(b)** Examples of colony phenotypes smooth and mucoid switch phenotypes originating from *C. neoformans* RC-2. F4-WR from 24067a and wrinkled and smooth from serotype A strain SB4. The colony morphologies were assessed after biofilms were disrupted and yeast cells were cultured on Sabouraud agar, and incubated for 48 h at 30°C




Fig. 2 *C. neoformans* phenotype affects plastic surface adhesion. Fungal cells adhesion to polystyrene was determined by cell counts. Results are representative of two experiments. Bars are the averages of three measurements, and error bars denote standard deviations. Asterisks denote *P* value significance calculated by *t*-test. *P* values of <0.05 were considered significant

switching frequency of biofilm-associated and planktonic cells of seven different variants which are derived from two different serotypes. Our data demonstrated that cells derived from biofilms and planktonic cells exhibited significant differences in switching rates. Biofilm-associated cells switched

more frequently than planktonic-grown cells in most strains and regardless of the phenotype. Although the increase was statistically significant the enhanced switch rates remained low (<10%). Four variants exhibited significantly more switches in their biofilm-associated cells when compared with their planktonic counterparts. In fact, there were a few variants that underwent multiple phenotypic switches. For example, WRS/SM-SM variant switched to WR, small, serrated, and sectorial colony phenotypes. In contrast, variants RC-2 MC and BF SB-4 WR exhibited significantly more reversion (switch back) to their smooth parent morphology as planktonic cells rather than as biofilm-associated cells.

Discussion

Phenotypic switching has been linked to the virulence of many pathogens, including fungi. High-frequency phenotypic switching is a phenomenon widely investigated for planktonic yeasts, including *C. neoformans* [4]. *C. neoformans* can undergo phenotypic switching in vitro to colony types that differ in their virulence in mice [14]. It is reasonable to speculate that phenotypic switching can also occur during cryptococcal biofilm formation. Therefore, the characteristics of *C. neoformans* phenotypic switching during biofilm formation were investigated. These phenotypes show variations in their capsular polysaccharide [14–16].

Table 1 Frequency in phenotypic switching of biofilm and planktonic cells of different phenotypes of *C. neoformans*

	Mucoid	Wrinkled	Small	Serrated	Sectored	Smooth
BF ^a RC-2-SM	(0.028%)		(1.1%)*			
PL ^a RC-2-SM	(0.05%)		(0.38%)			
BF WRS-SM		(0.6%)*	(3.0%)	(0.73%)*	(9.6%)*	
PL WRS-SM		(0.18%)	(9.6%)*	(0.25%)	(1.1%)	
BF F4-SM		(0.0015%)				
PL F4-SM		(0.0083%)				
BF SB-4-SM			(8.1%)*	(0.0021%)	(0.0021%)	
PL SB-4-SM			(5.2%)	(0.0016%)		
BF F4-WR			(1.4)*			(2.2%)*
PL F4-WR			(0.94)			(1.52%)
BF SB-4-WR						
PL SB-4-WR					(9.3%)	(46%)*
BF RC-2-MC						(0.004%)
PL RC-2-MC						(3.1%)*

^a BF and PL denote biofilm and planktonic cells, respectively

* Denotes *P* value significance calculated by *t*-test. *P* values of <0.05 were considered significant

Our data showed that phenotypic switch variants exhibited varying degrees of adhesion and biofilm production, indicating that phenotypic switching could be a critical factor affecting cryptococcal biofilm formation. In contrast to *Candida albicans* and *Candida parapsilosis* [17, 18], the smooth phenotype of *C. neoformans* exhibited the highest degree of both adhesion and biofilm formation. Biofilm formation was significantly higher than biofilm formation of mucoid and wrinkled switch variants of *C. neoformans*, suggesting that its superior biofilm-forming capability is at least in part a result of its higher ability to adhere to polystyrene. This observation was unexpected because wrinkled and mucoid switch variants typically exhibit larger polysaccharide capsules than their smooth parent counterparts.

The disparate effects of the capsule during biofilm formation may reflect differences in the chemical properties of the polysaccharide. Consistent with this view are the findings that *O*-acetylation of alginate in *Pseudomonas aeruginosa* contributed to biofilm formation [19] and the *C. neoformans* strain-related differences [10]. Previous work on the polysaccharide capsule of switch variants has shown that phenotypic switching can alter not only capsule size but also both the biochemical as well as the biophysical properties of the capsular polysaccharide [14, 15, 17]. These

changes can involve the position of the glucuronic acid on the mannose backbone as well as the distribution of charged residues over the mannose backbone. Although the molecular basis of this phenomenon has not been explored, the ability to bind polystyrene appeared to be inversely proportional to the complexity, and degree of substitution, of the GXM triad structure. Hence, the observation that the smooth phenotype made more robust biofilms combined with the phenotypic differences in binding polystyrene provided a compelling case for the importance of GXM binding as a critical event in *C. neoformans* biofilm formation.

In addition, our data confirm that the original colony phenotype (the one plated) dominates in biofilm, hence the difference in biofilm formation between switch variants could not be explained by differences in switching rate in biofilms. However our data also demonstrated that cells derived from cryptococcal biofilms and planktonic cells differed in phenotypic switch rate. These seemingly minor increases in switching rates in biofilms could however greatly affect the in vivo switching rate as the accumulation of biofilm-associated cells is likely to increase during chronic infection and switch variants are known to be selected in chronic infection [14] especially in the setting of therapeutic interventions

[6]. It is conceivable that, the smooth phenotype is more efficient at adherence and invasion but once it grows in cryptococcomas, which are biofilm-like associations in vivo, it can alter its phenotype to facilitate persistence. This may also explain why more diverse phenotypes are often observed in persistent infections [6].

Taken together our data indicate that different phenotypes of *C. neoformans* may be able to exploit different environmental niches and aggregate in biofilms in which they manifest altered switch rates. This links together two independent mechanisms that promote persistence during chronic infection and extends our understanding of fungal biofilm biology. Future studies might focus on delineating the environmental clues in biofilms that promote phenotypic switching.

Acknowledgment This work was supported by RO1 AI059681-04 grant and the Einstein's Hispanic Center of Excellence (HCOE).

References

- Mitchell TG, Perfect JR. Cryptococcosis in the era of AIDS—100 years after the discovery of *Cryptococcus neoformans*. Clin Microbiol Rev. 1995;8:515–48.
- Vecchiarelli A. Immunoregulation by capsular components of *Cryptococcus neoformans*. Med Mycol. 2000;38:407–17. doi:10.1080/714030973.
- Levitz SM. The ecology of *Cryptococcus neoformans* and the epidemiology of cryptococcosis. Rev Infect Dis. 1991;13:1163–69.
- Goldman D, Fries B, Franzot S, Montella L, Casadevall A. Phenotypic switching in the human pathogenic fungus *Cryptococcus neoformans* is associated with changes in virulence and pulmonary inflammatory response in rodents. Proc Natl Acad Sci USA. 1998;95:14967–72. doi:10.1073/pnas.95.25.14967.
- Guerrero A, Jain N, Goldman DL, Fries BC. Phenotypic switching in *Cryptococcus neoformans*. Microbiology. 2006;152:3–9. doi:10.1099/mic.0.28451-0.
- Fries BC, Cook E, Wang X, Casadevall A. Effects of antifungal interventions on the outcome of experimental infections with phenotypic switch variants of *Cryptococcus neoformans*. Antimicrob Agents Chemother. 2005;49:350–7. doi:10.1128/AAC.49.1.350-357.2005.
- Dromer F, Mathoulin-Pélissier S, Launay O, Lortholary O, The French Cryptococcosis Study Group. Determinants of disease presentation and outcome during cryptococcosis: The Crypto A/D Study. PLoS Medicine. 2007;4(2):e21. doi:10.1371/journal.pmed.0040021.
- Martinez LR, Casadevall A. *Cryptococcus neoformans* cells in biofilms are less susceptible than planktonic cells to antimicrobial molecules produced by the innate immune system. Infect Immun. 2006;74:6118–23. doi:10.1128/IAI.00995-06.
- Martinez LR, Casadevall A. Susceptibility of *Cryptococcus neoformans* biofilms to antifungal agents in vitro. Antimicrob Agents Chemother. 2006;50:1021–33. doi:10.1128/AAC.50.3.1021-1033.2006.
- Martinez LR, Casadevall A. Specific antibody can prevent fungal biofilm formation and this effect correlates with protective efficacy. Infect Immun. 2005;73:6350–62. doi:10.1128/IAI.73.10.6350-6362.2005.
- Walsh TJ, Schlegel R, Moody MM, Costerton JW, Salzman M. Ventriculoatrial shunt infection due to *Cryptococcus neoformans*: an ultrastructural and quantitative microbiological study. Neurosurgery. 1986;18:373–5. doi:10.1097/00006123-198603000-00025.
- Franzot SP, Mukherjee J, Cherniak R, Chen L, Hamdan JS, Casadevall A. Microevolution of a standard strain of *Cryptococcus neoformans* resulting in differences in virulence and other phenotypes. Infect Immun. 1998;66:89–97.
- Meshulam T, Levitz SM, Christin L, Diamond RD. A simplified new assay for assessment of fungal cell damage with the tetrazolium dye, (2,3)-bis-(2-methoxy-4-nitro-5-sulphenyl)-(2H)-tetrazolium-5-carboxanilide (XTT). J Infect Dis. 1995;172:1153–6.
- Fries BC, Taborda CP, Serfass E, Casadevall A. Phenotypic switching of *Cryptococcus neoformans* occurs in vivo and influences the outcome of infection. J Clin Invest. 2001;108:1639–48.
- McFadden DC, Fries BC, Wang F, Casadevall A. Capsule structural heterogeneity and antigenic variation in *Cryptococcus neoformans*. Eukaryot Cell. 2007;6:1464–73. doi:10.1128/EC.00162-07.
- Jin Y, Samaranayake YH, Yip HK, Samaranayake LP. Characterization of switch phenotypes in *Candida albicans* biofilms. Mycopathologia. 2005;160:191–200. doi:10.1007/s11046-005-6331-x.
- Fries BC, Goldman DL, Cherniak R, Ju R, Casadevall A. Phenotypic switching in *Cryptococcus neoformans* results in changes in cellular morphology and glucuronoxylomannan structure. Infect Immun. 1999;67:6076–83.
- Laffey SF, Butler G. Phenotype switching affects biofilm formation by *Candida parapsilosis*. Microbiology. 2005;151:1073–81. doi:10.1099/mic.0.27739-0.
- Nivens DE, Ohman DE, Williams J, Franklin MJ. Role of alginate and its O acetylation in formation of *Pseudomonas aeruginosa* microcolonies and biofilms. J Bacteriol. 2001;183:1047–57. doi:10.1128/JB.183.3.1047-1057.2001.